

Anti-MAPK1 Antibody, Rabbit Polyclonal

Synonyms: MAPK1, P38, P40, ERK2, ERT1, MAPK2, P42MAPK, P42-MAPK

Basic Info

Catalog

CP00004HuA20

Host

Rabbit

Conjugate

None

Size

100 μ L

Concentration

0.5 mg/mL

Physical State

Liquid

Species Reactivity

Human

Immunogen

Recombinant human MAPK1 protein,
fragment Val104~Lys342; UniprotKB:

P28482

Purification

Antigen Affinity Chromatography

Applications

WB

Property

Form & Buffer: Supplied in PBS, 50% glycerol, PH7.4.

Specificity / Sensitivity: Anti-MAPK1 Antibody, Rabbit Polyclonal recognizes endogenous levels of total MAPK1 protein.

Usage and Storage

Shipped at 4°C.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Avoid repeated freezing/thawing and violent shaking.

Please centrifuge it, before using.

Applications

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB: 0.5~5ug/mL

IF: 5~20ug/mL

IHC: 5~20ug/mL

ICC: 5~20ug/mL

QC Data




Figure 1. Application in WB

Western blot analysis of extracts of various cell lines and tissues, using MAPK1 antibody (CP00004HuA20) at 1 μ g/mL. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 8 s.

For Research Use Only!