

Anti-CKM Antibody, Rabbit Polyclonal

Synonyms: CK-MM, CKMM, CK3, CK-M, MM-CK, M-CK

Basic Info

Catalog

PA00011HuA10

Host

Rabbit

Conjugate

None

Size

100 μ L

Concentration

0.4 mg/mL

Physical State

Liquid

Species Reactivity

Predict reacts with: Human/Mouse/Rat

Immunogen

Recombinant human CKM protein,
fragment Met1~Lys381; UniprotKB:
P06732

Purification

Antigen Affinity Chromatography

Applications

WB

Property

Form & Buffer: Supplied in PBS, 50% glycerol, PH7.4.

Specificity / Sensitivity: Anti-CKM Antibody, Rabbit Polyclonal
recognizes endogenous levels of total CKM protein.

Usage and Storage

Shipped at 4°C.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Avoid repeated freezing/thawing and violent shaking.

Please centrifuge it, before using.

Applications

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB: 0.5~5 μ g/mL

IF: 5~20 μ g/mL

IHC: 5~20 μ g/mL

ICC: 5~20 μ g/mL

QC Data




Figure 1. Application in WB

Western blot analysis of extracts of various tissues, using CKM antibody (PA00011HuA10) at 1 μ g/mL dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 0.8s.

For Research Use Only!